

# Exotic Pest Fact Sheet 13

## Red Spider Mite (*Tetranychus evansi*)

### What is it?

Red spider mites prefer solanaceous crops and affect tomatoes but also eggplant, capsicums, potatoes and beans. Its small size makes it difficult to detect and it could easily be confused with other spider mites which may already be present in New Zealand.

### What does it look like?

Adult female Red spider mites are 0.5 mm long, oval-shaped, orange red coloured with an indistinct dark blotch on each side of the body (Fig. 1). They can lay up to 200 eggs. Males are smaller and straw to orange coloured. Their development is favoured by hot dry conditions and their life cycle is completed in 13.5 days.

Damage is mainly to the leaves of host plants. Feeding punctures created by the mites lead to whitening or yellowing of leaves, followed by desiccation, and eventually defoliation. In cases of severe attack, plants may die. An indication of Red spider mite is the presence of webbing which can be seen on the underside of the leaf (Fig. 2, Fig. 3), and in severe cases envelops parts of the plant.

### Why is it important?

Red spider mites infest the leaves of host plants and cause loss of yield. The pest does not directly affect fruit quality except for heavy infestations. Eradication can be difficult. Red spider mite is listed as a priority pest of concern by Biosecurity New Zealand.

### How does it spread?

Over short distances, mites can be spread by wind, irrigation water, and on clothing and tools. Mites can disperse within plants by walking. Movement of host plants can cause long distance spread. Its small size and similarity with other spider mite species makes it difficult to detect on consignments. Mites can be spread on the fruit of host plants, including tomatoes "on the vine", on Solanaceae plants, and as a hitchhiker on non-solanaceous plants.

### Where is it present?

Red spider mites originate from South America and were accidentally introduced to other parts of the world. It is now present in North and Central America, Caribbean, Southern and North Africa, France, Spain, Portugal, Italy, Greece, Japan, Taiwan, and in Australia where it is widespread in NSW and has been detected in Queensland on tomato plants.

### How can I protect my industry?

Check your production site frequently for the presence of new diseases and unusual symptoms. Make sure you are familiar with common pests and diseases of your industry so you can recognise something different.



**Fig 1:** Male (top) and female (bottom) Red spider mite. Image: Alain Migeon, CBGP – INRA, Monferrier-sur-Lez, France.



**Fig 2:** Red spider mite infested fruit. Image: Alain Migeon, CBGP – INRA, Monferrier-sur-Lez, France.



**Fig 3:** Red spider mite webbing around a leaf. Image: Alain Migeon, CBGP – INRA, Monferrier-sur-Lez, France.