

**Code of Practice  
for the Management of  
Potato Spindle Tuber Viroid (PSTVd)  
in Greenhouse Crops**

**February 2023**

## Table of Contents

1	Acknowledgements.....	4
2	Purpose of this Code of Practice .....	4
3	Scope.....	4
4	Benefits to Growers .....	5
5	Information on PSTVd.....	5
5.1	What is PSTVd? .....	5
5.2	Status of PSTVd in New Zealand .....	5
5.3	PSTVd symptoms.....	6
5.4	PSTVd transmission.....	7
5.5	Survival of PSTVd on surfaces .....	7
5.6	Disease prevention .....	7
6	Requirements of the Code of Practice.....	8
6.1	Responsibility for implementing the PSTVd code of practice.....	8
6.2	Source of planting material.....	8
6.3	Hygiene measures to prevent the spread of PSTVd .....	8
6.4	Surveying crops for PSTVd .....	9
6.5	Isolation of suspect PSTVd infected plants.....	9
6.6	Collection and submission of samples.....	9
6.7	Management of growing operations in greenhouses with suspect or confirmed PSTVd .....	9
6.8	Secure removal and destruction of confirmed PSTVd infected plants.....	9
6.9	Cleaning and disinfection between crop cycles.....	10
6.9.1	Greenhouse units with confirmed infections during the growth cycle .....	10
6.9.2	Greenhouses with no confirmed PSTVd during the growth cycle .....	10
6.10	Monitoring and record keeping .....	10
7	Compliance monitoring.....	10
8	Appendix 1. Procedure for compliance with the code of practice .....	11
8.1	Responsibility for compliance with this procedure .....	11
8.2	Source of planting material.....	11
8.3	Hygiene measures to prevent the introduction and spread of PSTVd .....	11
8.4	Surveying of crops for PSTVd .....	13
8.5	Isolation of suspect PSTVd infected plants.....	13
8.6	Collection and submission of samples .....	14
8.7	Management of growing operations in greenhouses with suspect or confirmed PSTVd ....	15
8.8	Secure removal and destruction of confirmed PSTVd infected plants.....	15
8.9	Disinfestation of the production unit at the end of the production cycle.....	15

8.10	Monitoring and record keeping .....	16
9	Appendix 2. Disinfectants .....	<b>Error! Bookmark not defined.</b>

## 1 Acknowledgements

This Code of Practice for the Management of Potato Spindle Tuber Viroid (PSTVd) in Greenhouse Crops is based on two previous documents:

- New Zealand Code of Practice for the Management of Potato Spindle Tuber Viroid (PSTVd) in Greenhouse Tomato and Capsicum Crops. New Zealand Vegetable & Potato Growers Federation, Wellington (Second Edition, 2003)
- Information on the Management of Potato Spindle Tuber Viroid (PSTVd) in Greenhouse Tomato and Capsicum Crops. Tomatoes New Zealand Inc. and Vegetables New Zealand Inc., Wellington (Draft, 2015).

The code of practice also includes procedures for destruction of infected material and cleaning and disinfection adapted from MPI's Operational Specification for the Potato Spindle Tuber Viroid (PSTVd) response (2022). The publishers of, and contributors to, those three documents are acknowledged.

## 2 Purpose of this Code of Practice

The code of practice supports the objectives of the Potato Spindle Tuber Viroid (PSTVd) 2022 Response Operational Agreement, which are:

- To rapidly detect PSTVd and delimit its spread within New Zealand
- To determine the extent of spread on the infected property
- To eradicate the viroid from within New Zealand
- To minimise impacts to potentially impacted industries

The 2022 PSTVd response has aimed to rapidly eradicate PSTVd by destruction of affected crops, cleaning and disinfection, and greenhouse hygiene. This approach has been constrained by several factors:

- The strain of PSTVd detected shows mild symptoms, making detection difficult.
- The time from infection to plants displaying symptoms can be several weeks or months, presenting the risk of transmission from symptomless plants.
- A mandatory programme involving all growers is not feasible.
- Crop destruction is constrained by cost.

The code of practice may be adopted by Response Governance as an operational procedure for eradication of PSTVd through grower management of the crop through to the end of a crop cycle, with oversight from MPI's Response Operations.

The code of practice is recommended practice for other commercial growers to rapidly detect and contain the spread of PSTVd within greenhouses or to other crops (other greenhouses, properties, or species – especially potatoes).

## 3 Scope

This code of practice applies to commercial greenhouse production tomatoes, capsicums, and eggplants. The requirements apply in two situations:

- In greenhouses where PSTVd has been confirmed and greenhouse operators are directed by MPI to operate in accordance with this code of practice; and

- In all other commercial greenhouses as a recommended practice for greenhouse hygiene, disease prevention, and early PSTVd detection.

## 4 Benefits to Growers

Implementation of this code of practice has benefits to individual growers and to the industry overall. The direct benefits are:

- The overall direct cost of the response will be lower.
- Impacts on affected growers will be lower, and therefore more certainty regarding ongoing production and sale of fruit from affected properties.
- Potential production losses from PSTVd will be minimised.
- Good hygiene practices will also reduce the potential spread of other diseases within greenhouse properties.
- The programme will maintain market access as the status of PSTVd will be maintained as Transient: Under Eradication.
- Potential for early detection of other exotic diseases.
- Reduced risk of establishment of PSTVd in other hosts e.g., solanaceous weeds and ornamentals, or into potato production.

## 5 Information on PSTVd

### 5.1 What is PSTVd?

Potato spindle tuber viroid (PSTVd) was first reported in the USA in 1922 and, as its name suggests, causes spindle shaped potato tubers. Since that time it has spread to many potato producing countries. PSTVd also infects other solanaceous crops including tomato, capsicum, aubergine, pepino, and cape gooseberry as well as wild and ornamental solanaceous species (e.g., nightshade, Poroporo, jasmine nightshade).

Viroids are smaller than viruses, but replicate in a similar way. They take over host cells within a plant and uses the plant's metabolic process to produce more viroid particles. The infection progresses to a stage where the metabolism in the whole plant is affected. With PSTVd the time from infection to expression of symptoms can take several weeks. During that time the plant may be infectious and presents a risk to transmission between plants.

### 5.2 Status of PSTVd in New Zealand

The official status of PSTVd in New Zealand is:

*Transient: actionable, under eradication. PSTVd has been detected as an isolated population which may survive into the immediate future and, without phytosanitary measures for eradication, may establish. Appropriate phytosanitary measures have been applied for its eradication.*

As noted in Section 2, this code of practice forms part of those measures being applied to eradicate PSTVd from New Zealand. PSTVd is a regulated pest for New Zealand and designated as an "unwanted organism" under the Biosecurity Act 1993.

PSTVd has been detected on a few occasions in New Zealand on greenhouse crops and one instance on cape gooseberry, but has been eradicated each time. PSTVd has never been detected in New Zealand on potato, and there are no records of PSTVd in outdoor tomato or capsicum crops in New Zealand. There are also no records of detection of the viroid from wild host plants in New Zealand.

It is thought that previous incursions of PSTVd have been carried on imported seed.

### 5.3 PSTVd symptoms

PSTVd is difficult to detect and may be present without any visible signs of infection showing in a plant. Symptoms may take several weeks from infection before symptoms become visible. This presents a risk of transmission to other plants.

Visual signs of PSTVd can be similar to those caused by nutrient deficiency, root disease and pest or spray drift damage.

#### 5.3.1 Tomato symptoms

In tomatoes, there is a long latent period of four to five weeks after infection before the symptoms are expressed. Overseas, young plants (24 true leaf stage) infected with the virulent strain have shown rugosity, epinasty and curling in new leaflets. The internodes become shorter, making the plant appear stunted and malnourished. However, symptoms are not usually evident until after the 5th truss. Later, severe veinal necrosis may develop. So far no fruit symptoms have been described in tomatoes other than discolouration (yellowing) and blotchiness. Symptoms are exhibited more strongly under high radiation conditions suggesting that spring and summer crops may be more susceptible.

Yield of tomatoes from affected plants will be reduced, however losses depend on the age of the plant and the severity and strain of the infection. Lost production may be from 10-20%, and as high as 60% with severe strains.

Further information can be found in the [Tomatoes NZ factsheet](#) or the [Horticulture NZ factsheet](#).

#### 5.3.2 Capsicum symptoms

In capsicums, PSTVd can show itself in very subtle ways. Some common indicators are:

- Ruffled or wavy edges to the leaves. (Note: some varieties; e.g. the orange variety Boogie, have naturally ruffled or wavy leaves.)
- Leaves may be smaller than normal.
- The calyx may curl up at the edges.
- Fruit may be smaller than normal.
- The fruit stem may be thin and skinny looking.
- Plant heads tend to be smaller and thinner than normal.

The severity of PSTVd in capsicums may vary from crop to crop and from variety to variety, as some varieties may be more susceptible to PSTVd infection than others. The production of capsicums from affected plants may not be significantly reduced, however the fruit is normally smaller and the calyx may curl up at the edges.

### 5.3.3 Aubergine / eggplant symptoms

### 5.3.4 Pepino symptoms

## 5.4 PSTVd transmission

PSTVd is transmitted in four different ways:

- Mechanical transmission. The main way that PSTVd is transmitted within crops is through routine plant handling operations and transfer of PSTVd on knives, equipment and clothing. Leaf hairs contain high concentrations of viroid and are easily transferred to other surfaces by normal cultivation activities. This occurs with potatoes and tomatoes, where viroid spread is mainly along the row.
- Infected seed and pollen. PSTVd is transmitted on seed and also on pollen. The rate of seed transmission is variable and can depend on variety. All solanaceous seed is imported to New Zealand under import conditions with strict requirements for PSTVd freedom.
- Aphid transmission. PSTVd can be transmitted by *Myzus persicae* (peach potato aphid) when infected with both *Potato leafroll virus* (PLRV) and PSTVd, thereby limiting the number of potential infection sources.
- Vegetative propagation. PSTVd can be transmitted from infected material during grafting and propagation by cuttings.

## 5.5 Survival of PSTVd on surfaces

PSTVd can survive in dried tomato sap for more than eight weeks and in infected leaf debris for over six months (DPI-NSW, 2023<sup>1</sup>).

PSTVd survives on many contaminated materials, remaining infective on contaminated metal after 1 hour; contaminated cotton after 6 hours; and contaminated wood, plastic and leather remained infective after 24 hours (Barbetti, 2012)<sup>2</sup>. However PSTVd survived for only 30 minutes on human skin. PSTVd survived for at least 4 weeks on aluminium but survived poorly on plastic (Spence and O'Neill, 2004<sup>3</sup>).

## 5.6 Disease prevention

Prevention includes all measures to prevent the introduction of PSTVd into a crop. It is very important to start a new production cycle with viroid-free planting material (seeds or plants).

In addition to the use of healthy planting material, it is also important to prevent viroid introduction via human activities. Because PSTVd is easily transmitted mechanically it can be introduced into a greenhouse from other infected sources via the hands, clothes, or equipment used by people working in or visiting the

---

<sup>1</sup> <https://www.dpi.nsw.gov.au/biosecurity/plant/insect-pests-and-plant-diseases/pstv>

<sup>2</sup> [https://ausveg.com.au/app/data/technical-insights/docs/130014\\_VG09110.pdf](https://ausveg.com.au/app/data/technical-insights/docs/130014_VG09110.pdf)

<sup>3</sup> Protected tomato: Sources, survival, spread and disinfection of Potato spindle tuber viroid (PSTVd). Final report to Horticultural Development Council, UK. 2004

greenhouse. The use of disposable gloves and specific clothing and equipment that stays inside a greenhouse unit may help prevent PSTVd spread to other greenhouse units.

Eradication is based on destruction of PSTVd-infected plants and thorough cleaning of equipment and greenhouses where infected plants have been grown.

## 6 Requirements of the Code of Practice

Key requirements of the code of practice are:

- Hygiene measures to prevent spread of PSTVd into and within greenhouses
- Surveying of greenhouse crops for PSTVd
- Isolation of suspect PSTVd infected plants and sampling for testing
- Secure removal and destruction of confirmed PSTVd infected plants
- Management of growing operations in greenhouses with suspect or confirmed PSTVd
- Cleaning and disinfection between growing cycles.

These requirements are presented below, and growers should consider how they will meet the requirements. PSTVd is highly transmissible through handling and working with plants. Ideally, the requirements will fit in with existing hygiene practices, but growers should review their hygiene and isolation practices against these requirements and modify practices if necessary.

Growers should document how they will meet the requirements, so that all staff are aware of their roles in PSTVd management. The procedures need to identify what actions are taken, who is responsible for the action, when it will be done, and how the outcome will be recorded. Alternatively, growers can adopt or modify the generic procedure provided in Appendix 1.

### 6.1 Responsibility for implementing the PSTVd code of practice

Greenhouse operators should nominate a person responsible for implementing the code of practice including monitoring of hygiene procedures, surveying of crops, sampling and isolation suspected PSTVd infected plants, removal and destruction of confirmed PSTVd infected plants, management of harvest operations, supervision of disinfection of production sites, and record keeping. This person is referred to as the PSTVd manager.

### 6.2 Source of planting material

Greenhouse operators should ensure that planting material is free from PSTVd. This can be achieved by ensuring that seed or plant suppliers have had the seed tested and found free from PSTVd, and are part of the Plant Pass scheme.

### 6.3 Hygiene measures to prevent the spread of PSTVd

The activities and movement of people and equipment is one of the main ways that PSTVd is spread within and between greenhouses. Greenhouse operators or PSTVd managers will document the procedures that they will follow to minimise the likelihood of PSTVd entering greenhouses and spreading between plants and between units or other production sites. These procedures must address the following:

- Ensuring the greenhouse environs are free from non-crop PSTVd host plants;
- control over movement of staff and equipment (including pollinators) into and between greenhouse units ;
- how hands are kept clean;
- how knives and tools are kept clean or sterilised;
- cleaning of equipment such as trolleys;
- control over visitors to the greenhouse;
- removal and destruction of volunteer plants.

#### 6.4 Surveying crops for PSTVd

Greenhouse operators or PSTVd managers will document procedures for monitoring greenhouses for PSTVd. The monitoring procedures must:

- Identify who will undertake surveying activities;
- Describe how surveying will be undertaken;
- Ensure that the surveying will be carried out at least
  - weekly in infected greenhouses
  - At least monthly in all other greenhouses;
- Describe how proficiency of persons surveying for PSTVd will be ensured;
- Describe how samples of plant material from suspected PSTVd infected plants will be taken and submitted for identification;
- Describe how records will be kept.

#### 6.5 Isolation of suspect PSTVd infected plants

Greenhouse operators or PSTVd managers will document procedures for dealing with suspect and confirmed PSTVd infections. The procedures must:

- Describe how the suspect plant and the 5 plants on either side will be marked to allow for future identification;
- Describe how the suspect plant will be isolated from the rest of the production unit (to prevent possible spread of PSTVd);
- Describe how the marked plants will be isolated and additional hygiene measures applied to prevent disease spread until test results are received.

#### 6.6 Collection and submission of samples

Where it is suspected that a plant is infected with PSTVd, take leaf samples and submit them to the MPI Approved Laboratory in accordance with Appendix 1 (8.6 Collection and submission of samples).

#### 6.7 Management of growing operations in greenhouses with confirmed PSTVd

To be updated

#### 6.8 Secure removal and destruction of confirmed PSTVd infected plants

- Describe the procedure for removal of a confirmed PSTVd infected plant and the 5 plants on either side ensuring:
  - staff wear disposable gloves
  - plants are bagged for disposal
  - plants are burned or deep -buried
  - equipment used with the infected plants is sterilised or disposed of.

## 6.9 Cleaning and disinfection between crop cycles

PSTVd can be difficult to detect and may be present in a greenhouse without the grower's knowledge. Thorough cleaning and disinfection between crop cycles is an opportunity to break a disease cycle by removing potentially infected plants and cleaning surfaces where PSTVd may survive and be transferred to new plants. Cleaning and disinfection practices differ depending on whether PSTVd has been detected previously in a greenhouse unit.

### 6.9.1 Greenhouse units with confirmed infections during the growth cycle

Greenhouse units where there have been confirmed PSTVd infections must follow the cleaning, disinfection and disposal procedure described in Appendix 1, Section 8.9. This includes secure removal of plant material for deep burial, as well as cleaning and disinfection of the greenhouse unit under MPI supervision.

### 6.9.2 All other greenhouses (no confirmed PSTVd during the growth cycle)

Despite there having been no confirmed detections of PSTVd, there may be latent or undetected PSTVd infection present. To break any potential disease cycle growers should follow the procedure described in Appendix two, however disposal by deep burial is not a requirement.

## 6.10 Monitoring and record keeping

The PSTVd manager will monitor CoP activities and keep records of:

- Surveying, including the location and marking of suspect infections
- Locations and dates of samples submitted for diagnosis, and the results
- Actions taken on receipt of PSTVd positive results
- disinfection operations between crop cycles.

## 7 Compliance monitoring

To be updated

## 8 Appendix 1. Procedure for compliance with the code of practice

### 8.1 Responsibility for compliance with this procedure

The PSTVd manager is responsible for this procedure. The PSTVd manager for [Business name and address] is [Name.....]

### 8.2 Source of planting material

- Record details of planting material, such as
  - Source of seeds (including batch numbers, supplier, dates) used for rootstocks and grafting
  - Confirmation by suppliers that seeds
    - were imported in compliance with MPI's Import Health Standard, or
    - if New Zealand grown a representative sample of at least 3000 seeds, drawn according to the ISTA or AOSA sampling methodology, was tested and found free from PSTVd
  - Dates of receipt and planting (including units / rows planted)

### 8.3 Hygiene measures to prevent the introduction and spread of PSTVd

- Ensure greenhouse environs are free from wild and ornamental hosts of PSTVd
  - Inspect the area around the greenhouse and remove any wild host plants such as nightshade (refer HOST LIST link)
- Ensure staff are familiar with requirements for cleaning and disinfection, including use of disinfectant products:
  - General equipment to be cleaned and disinfected with virkon at 2% with contact time of at least 1 hour, rinse with water afterwards.
  - Sensitive equipment cleaned with virkon 2% should then be wiped with 70% ethanol to remove any residual virkon
  - Scissor trolleys – make watertight and steam clean. Wipe external parts of the trolley with virkon 2% where it is safe and practical to do so. Rinse with water after.
  - Equipment able to be treated with virkon 2% must follow the label directions for use, with an increased contact time to 1 hour. Rinse with water afterwards.
- Develop and follow a containment plan for the greenhouse complex (separating the greenhouse complex into units where movement of people and equipment between units is minimised may reduce the spread of PSTVd across the site)
  - Identify all greenhouses on a diagram
  - Identify which greenhouses (or sections of greenhouses) are considered separate growing units because -
    - They have doors between sections
    - Are managed by separate teams or on separate days
    - Have equipment (e.g., trolleys, knives) that stay within the unit and are not shared between units
    - Have independent hydroponic systems
  - Instruct all staff on the operation of the containment plan and monitor implementation.

- Instruct all staff on hygiene requirements and monitor implementation:
  - No animals are allowed inside the greenhouse at any time (other than pollinators);
  - Staff must wash and sanitise hands when entering or leaving the unit and before and after breaks;
  - Staff working the crop must
    - wear latex (or other impermeable) gloves that are disposed of at the end of each day or when moving to a new unit;
    - Dip gloved fingers and knives into disinfectant solution before moving into the crop or before commencing work in a new unit;
    - Dip gloved fingers or pruning tools into disinfectant solution inside a container and/or sponge when twisting, layering or leafing;
    - Replace disinfectant frequently (as these products break down in contact with light and organic matter) and wash sponges daily;
    - Perform all plant handling tasks such as twisting, leaf removal and picking in the same direction along the row. (This is very important, as this will ensure that if PSTVd does spread in the greenhouse, it will spread mainly in one direction).
- Instruct all visitors on hygiene requirements and monitor implementation
  - Contractors entering the premises to work on plants wear new gloves and clean protective clothing (or disposable suits) that remain in the greenhouse or are disposed of when finished;
  - Advisors, seed and other company representatives wear clean protective clothing and also gloves if handling plants;
- Supervise visitors not working with the crop to ensure that they keep to the pathways and do not handle the crop;
- Place mats at all doorways and keep them moist with disinfectant throughout the day.
- Ensure staff comply with equipment cleaning instructions:
  - Clean and spray trolleys with disinfectant plus any other equipment being moved from one unit to another;
  - Sweep and clean trolleys and bins before moving to a new row ;
  - Keep concrete paths free of plant material at all times by ensuring bins are not overfilled and any spillage is promptly cleaned up;
  - Place leaves and prunings into designated containers;
  - Don't get into bins to tramp down leaf matter;
  - Keep trolleys free of plant material and sweep and clean at the end of each day;
  - Sanitise reused equipment; e.g. bobbins, clips;
- Ensure that pollinator colonies are not moved between units.
- Ensure staff removing rogue plants:
  - Remove any tomato or capsicum seedlings emerging from the drains or holes;
  - Do this as a separate job and ensure other plant material is not touched;

- Wear gloves and dispose of these afterwards.

#### 8.4 Surveying of crops for PSTVd

- Ensure that all staff working plants in the designated production units exhibit competency in the recognition and reporting of symptoms of PSTVd.
  - Instruct staff on PSTVd symptoms and reporting of suspect PSTVd to the PSTVd surveyor
  - Ensure that PSTVd identification resources (e.g., booklets distributed by TNZ and VNZ) are readily available to all staff working the crop during the surveying period.
- Ensure that greenhouses are surveyed for PSTVd at least weekly in infected greenhouses and at least monthly in all other greenhouses
  - Survey activities may be part of normal plant management activities
  - Ensure that survey personnel are competent in recognition of PSTVd symptoms and recording and reporting requirements
    - Train survey personnel in....
    - Assess competency of survey personnel in symptom recognition, recording and reporting
    - Maintain records of competency assessments
- During the surveying programme, ensure that survey personnel inspect each plant for symptoms of PSTVd.
- Monitor survey personnel during PSTVd surveying to ensure that where symptoms resembling those of PSTVd are detected, these are reported to the PSTVd manager.
- Inspect suspect plants and determine whether the plants are likely to be infected with PSTVd, or if there is another cause of infection.
- Maintain records of surveying activities and actions taken.

#### 8.5 Isolation of suspect PSTVd infected plants

- When a plant has been sampled for suspect PSTVd, attach a label to the plant so that it can be identified as a suspect plant (in case resampling is necessary or if positive results are obtained).
- Attach a label to the 5 plants on either side of the suspect plant, indicating that they are adjacent to a suspect plant.
- Cordon off the row (both sides) and ensure that staff working in the area are aware of suspect PSTVd plants in that row and of the following requirements.
- Ensure staff working plants in the rows with suspected infected plants follow additional hygiene measures:
  - Work the isolated row from each end if possible, working towards the isolated, working the isolated plants last.
  - wear latex (or other impermeable) gloves that are disposed of at the end of each day or when moving to a new row;
  - Dip gloved fingers and knives into disinfectant solution before moving into the crop or before commencing work in a new row;
  - Dip gloved fingers or pruning tools into disinfectant solution inside a container and/or sponge when twisting, layering or leafing;
  - Replace disinfectant frequently (as these products break down in contact with light and organic matter) and wash sponges daily;

- Clean and spray trolleys with disinfectant plus any other equipment being moved from one row to another;
- Place leaves and prunings into designated containers;
- Don't get into bins to tramp down leaf matter;
- Sweep and clean trolleys and bins before moving to a new row ;
- Keep concrete paths free of plant material at all times by ensuring bins are not overfilled and any spillage is promptly cleaned up;
- Keep trolleys free of plant material and sweep and clean at the end of working the affected row;
- Sanitise reused equipment; e.g. bobbins, clips;
- Fruit from suspect plants may be harvested and sold, unless the diagnosis confirms PSTVd.

### 8.6 Collection and submission of samples

Where it is suspected that a plant is infected with PSTVd, take leaf samples and submit them to the MPI Approved Laboratory in accordance with the following procedure:

- Do not touch the suspect plants with bare hands but wear disposable gloves for sampling.
- Use a new pair of disposable gloves for each plant sampled.
- Take a sample of 5 leaves from each suspect plant. Take leaves at equal distances from between 5cm from the top down to 1m from the top.
- Collect each sample of 5 leaves using gloved hand and while still holding sample turn glove inside out leaving sample well inside (so the leaves do not come out of the glove).
- Place samples in a ziplock plastic bag with the air expressed prior to sealing.
- This bag should be placed inside another ziplock bag and sealed.
- Label the bag with full details of the grower's name, address, and contact phone numbers.
- Keep bags containing leaf samples cool at all times and ensure they are not frozen. Do not leave samples in the sun or inside hot vehicles. Place samples in a small chilly bin with a cold pack wrapped in several paper towels or newspaper to prevent freezing of samples.
- Wash hands thoroughly with soap and water and decontaminate any equipment.
- Label the chilly bin with laboratory address and full grower return address and contact phone numbers.
- Contact the laboratory in advance of sending off samples to advise that a suspect sample is being submitted. If samples are collected on Sunday through Thursday, dispatch samples immediately on overnight courier. For samples collected on Friday or Saturday, leave a message on the answer phone and keep samples cool, and courier to the laboratory on Sunday (for Monday delivery).

Attention: Lisa Ward

URGENT- Virology – Plant sample for testing

MPI PHEL

231 Morrin Road

St Johns

AUCKLAND

Ph: 09 909 3015

Fax: 09 909 5739

Email: [lisa.ward@mpi.govt.nz](mailto:lisa.ward@mpi.govt.nz)

## 8.7 Management of growing operations in greenhouses with confirmed PSTVd

- New section – see 6.7

## 8.8 Secure removal and destruction of confirmed PSTVd infected plants

- If laboratory tests confirm PSTVd, supervise staff in the removal of suspect plants and the 10 marked neighbouring plants as follows:
  - Wear gloves and protective clothing when removing infected plants;
  - Remove plants and place in bags for disposal;
  - Dispose of plant material by burning or deep burial;
  - Dispose of growing media by deep burial;
  - Dispose of, or sanitise, other equipment (such as bobbins, support clips and truss clips) used in conjunction with infected plants.

## 8.9 Cleaning and disinfection between crop cycles

- At the end of production, supervise staff in the removal of all plants, their disposal, and cleaning and disinfection of surfaces in the production unit.
- Confirm all persons involved in this process wear disposable suits that are left in the greenhouse during breaks and discarded at the end of the cleanup.
- Ensure that plant material and growing slabs/bags are removed as quickly as possible and the floor is swept clean of all debris.
- Confirm that the plant material is transferred to vehicles and taken for disposal.
- After the crop has been removed confirm that the following is undertaken:
  - Waterblast gutters and troughs and floors.
  - Spray and treat concrete floors and paths with virkon 2% with contact time of 1 hour. Rinse with water afterwards.
  - Clean and disinfect plastic sheeting with Virkon or alternatively remove it dispose of;
  - Treated glass house walls with virkon 2%, then rinse with water. Only clean the glasshouse walls as high as the second windowpane taking care to avoid splashing any virkon onto the ceiling and screens. Then clean the remainder with GS4 before washing with water.
  - Clean the underside of the roof glass and ceiling screens using the underside roof washer with pressurised fresh water.
  - Flush and sterilize all irrigation lines with nitric acid (following the label directions for concentration) to remove build-up of organic material, and then flush with water. The system should then be filled with hyperstat (following the label directions for concentration) which will sit there for a week before it is then flushed with water again

- Replace CO2 lines;
- Replace or sanitise all bobbins, support clips and truss clips.
- Disposal of pollinator colonies.
- Water tank systems are decontaminated and cleaned out with sodium hypochlorite following the label directions for concentration, with a contact time of 1 hour. Flush thoroughly with water. Check with supplier/technician for compatibility.
- If biofilters are used, consult the supplier for decontamination procedures.
- Maintain record of disinfection operations, including the following information:
  - Date harvesting finished;
  - Date(s) disinfection commenced and finished;
  - Method and location of disposal of plant material;
  - Method of treatment/disposal of growing substrate and other substrate;
  - Chemicals used to disinfect/sanitise.

#### 8.10 Monitoring and record keeping